

MSUMB 96-02, October 1996

h -deformation of grassmann supergroup $Gr(1|1)$

Salih Çelik^{1 2}

Mimar Sinan University, Department of Mathematics,
80690 Besiktas, Istanbul, TURKEY.

Abstract

We introduce the h -deformation of the algebra of functions on the grassmann supergroup $Gr(1|1)$ via a contraction of $Gr_q(1|1)$.

¹E-mail: celik@msu.edu.tr

²New E-mail: sacelik@yildiz.edu.tr

Recently matrix groups like $GL(2)$, $GL(1|1)$, *etc.*, were generalized in two ways. Both are based on the deformation of the algebra of functions on the groups generated by coordinate functions t^i_j that commute

$$[t^i_j, t^k_l] = 0.$$

In the q -deformation of matrix groups [1], these commutation relations are determined by a matrix R_q so that the functions do not commute but satisfy the equation

$$R_q(T \otimes T) = (I \otimes T)(T \otimes I)R_q.$$

In this equation the elements of R_q are numbers but the matrix $T = (t^i_j)$ is formed by generally non-commuting elements of an algebra.

Another type of deformation, so called the " h -deformation", which is a new class of quantum deformations of Lie groups and Lie algebras has recently been intensively studied [2-4]. These deformations can be important since the deformation parameter h can directly be identified with the Planck constant, which makes them interesting for physical applications. The other important point is that the standard q -deformation and the h -deformation have been characterized as the only two deformations of $SL(2)$ [3].

The goal of this paper is to develop the h -deformation for the case of grassmann matrix supergroups. We study the simplest grassmann supergroup $Gr(1|1)$, here. The one parametric q -deformation of $Gr(1|1)$, $Gr_q(1|1)$, was given in Ref. 5. Before discussing the h -deformation of $Gr(1|1)$, we give some notations and useful formulas in following.

A grassmann supermatrix $\hat{T}$, that is, an element of $Gr(1|1)$, is of the form

$$\hat{T} = \begin{pmatrix} \alpha & b \\ c & \delta \end{pmatrix}$$

with two odd (greek letters) and two even (latin letters) matrix elements. The following closely follows the approach of Ref. 5.

In this paper we denote q -deformed objects by primed quantities. Unprimed quantities represent transformed coordinates.

Let us consider the following quantum superplane and its dual, denoted by A_q and A_q^* , respectively [6]:

$$U' = \begin{pmatrix} x' \\ \xi' \end{pmatrix} \in A_q \iff x'\xi' - q\xi'x' = 0, \quad \xi'^2 = 0, \quad (1)$$

and its dual

$$\hat{U}' = \begin{pmatrix} \eta' \\ y' \end{pmatrix} \in A_q^* \iff \eta'^2 = 0, \quad \eta' y' - q^{-1} y' \eta' = 0. \quad (2)$$

Suppose that the matrix elements of $\hat{T}'$ (anti-)commute with the coordinates of A_q and A_q^* . Then, the endomorphisms

$$\hat{T}' : A_q \longrightarrow A_q^* \quad \text{and} \quad \hat{T}' : A_q^* \longrightarrow A_q \quad (3)$$

impose the following bilinear product relations among the matrix elements of $\hat{T}'$:

$$\begin{aligned} \alpha' b' &= q^{-1} b' \alpha', & \alpha' c' &= q^{-1} c' \alpha', \\ \delta' b' &= q^{-1} b' \delta', & \delta' c' &= q^{-1} c' \delta', \\ \alpha' \delta' + \delta' \alpha' &= 0, & \alpha'^2 &= 0 = \delta'^2, \\ b' c' &= c' b' + (q - q^{-1}) \delta' \alpha' \end{aligned} \quad (4)$$

where q is a non-zero complex number and $q^2 \neq 1$. These relations are the defining relations of $\text{Gr}_q(1|1)$.

We introduce new coordinates x and ξ by

$$U = g^{-1} U' \quad (5)$$

where

$$g = \begin{pmatrix} 1 & 0 \\ h/(q-1) & 1 \end{pmatrix} \quad (6)$$

as in Ref. 7. Here the deformation parameter h is a grassmann number which has the following properties

$$h^2 = 0 \quad \text{and} \quad h\xi = -\xi h.$$

Now, in the limit $q \rightarrow 1$ we get the following exchange relations, which define the h -superplane A_h

$$x\xi = \xi x + hx^2, \quad \xi^2 = -hx\xi. \quad (7)$$

Similarly, using the relations (2) with (6), one has [after taking the limit $q \rightarrow 1$] the dual h -superplane A_h^* as generated by η and y with the relations

$$\eta^2 = 0, \quad \eta y = y\eta. \quad (8)$$

We now consider the endomorphisms

$$\hat{T} : A_h \longrightarrow A_h^* \quad \text{and} \quad \hat{T} : A_h^* \longrightarrow A_h. \quad (9)$$

Then, we define the corresponding h -deformation of the grassmann supergroup $\text{Gr}(1|1)$ as a quantum matrix supergroup $\text{Gr}_h(1|1)$ generated by α , b , c and δ which satisfy the following h -commutation relations

$$\begin{aligned} \alpha b &= b\alpha + hb^2, & \alpha c &= c\alpha + h(bc + \delta\alpha), \\ \delta b &= b\delta - hb^2, & \delta c &= c\delta - h(cb - \alpha\delta), \\ \alpha^2 &= h\alpha b, & \alpha\delta &= -\delta\alpha + h(\delta b - b\alpha), \\ \delta^2 &= -h\delta b, & bc &= cb + h(b\delta + \alpha b). \end{aligned} \quad (10)$$

It can be check that the maps

$$\mu : A_h \longrightarrow \text{Gr}(1|1) \otimes A_h^*, \quad \mu^* : A_h^* \longrightarrow \text{Gr}(1|1) \otimes A_h \quad (11)$$

such that

$$\begin{aligned} \mu(U) &= \hat{T} \otimes \hat{U}, \quad i.e. \quad \mu(u_i) = t_i^j \otimes \hat{u}_j, \\ \mu^*(\hat{U}) &= \hat{T} \otimes U, \quad i.e. \quad \mu^*(\hat{u}_i) = t_i^j \otimes u_j \end{aligned} \quad (12)$$

define the co-action of the quantum group $\text{Gr}_h(1|1)$ on the quantum superplane A_h and its dual A_h^* , respectively.

An interesting feature is that the second column of the matrix $\hat{T} \in \text{Gr}_h(1|1)$ is isomorphic to A_h , though this is not so in $\text{GL}_h(1|1)$ [7].

Alternatively, the relations (10) can be obtained using the following similarity transformation which given by Aghamohammadi *et al* [4]:

$$\hat{T}' = g\hat{T}g^{-1}. \quad (13)$$

To do this, we use the relations (4) and next the limit $q \rightarrow 1$.

The quantum (dual) superdeterminant of $\hat{T}$ is again defined as [5]

$$\hat{D}_h = bc^{-1} - \alpha c^{-1}\delta c^{-1} = c^{-1}b - c^{-1}\alpha c^{-1}\delta \quad (14)$$

which is independent of the relations (10). It is easy to verify that $\hat{D}_h$ commutes with all matrix elements of $\hat{T}$ (and h), that is, $\hat{D}_h$ belongs to the centre of the algebra

$$\hat{T}\hat{D} = \hat{D}\hat{T}.$$

The inverse of $\hat{T}$ is defined as [5]

$$\hat{T}^{-1} = \begin{pmatrix} -c^{-1}\delta b^{-1} & c^{-1} + c^{-1}\delta b^{-1}\alpha c^{-1} \\ b^{-1} + b^{-1}\alpha c^{-1}\delta b^{-1} & -b^{-1}\alpha c^{-1} \end{pmatrix} \quad (15)$$

provided b and c are invertible.

We shall now obtain an R-matrix for the quantum grassmann supergroup $\text{Gr}_h(1|1)$ from the R-matrix of $\text{Gr}_q(1|1)$ which is introduced in Ref. 8.

The associative algebra (4) is equivalent to

$$R_q \hat{T}'_1 \hat{T}'_2 = -\hat{T}'_2 \hat{T}'_1 R_q \quad (16)$$

where [8]

$$R_q = \begin{pmatrix} q + q^{-1} & 0 & 0 & 0 \\ 0 & -2 & q^{-1} - q & 0 \\ 0 & q - q^{-1} & -2 & 0 \\ 0 & 0 & 0 & q + q^{-1} \end{pmatrix}. \quad (17)$$

Here, we used the tensoring convention

$$\begin{aligned} (\hat{T}_1)^{ij}_{kl} &= (\hat{T} \otimes I)^{ij}_{kl} = (-1)^{k(j+l)} \hat{T}^i_k \delta^j_l, \\ (\hat{T}_2)^{ij}_{kl} &= (I \otimes \hat{T})^{ij}_{kl} = (-1)^{i(j+l)} \hat{T}^j_l \delta^i_k. \end{aligned} \quad (18)$$

Note that although the algebra (4) is an associative algebra of the matrix elements of $\hat{T}'$, R_q does not satisfy the (graded) quantum Yang-Baxter equation (QYBE)

$$R_{12} R_{13} R_{23} = R_{23} R_{13} R_{12}. \quad (19)$$

Thus the QYBE is not a necessary condition for associativity. However, as interesting point, if we decompose the matrix R_q is of the form

$$R_q = R_q^1 + R_q^2 \quad (20)$$

where

$$R_q^1 = \begin{pmatrix} q & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & q - q^{-1} & -1 & 0 \\ 0 & 0 & 0 & q^{-1} \end{pmatrix}, \quad R_q^2 = \begin{pmatrix} q^{-1} & 0 & 0 & 0 \\ 0 & -1 & q^{-1} - q & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & q \end{pmatrix} \quad (21)$$

then both matrices R_q^1 and R_q^2 do satisfy the (graded) QYBE. Thus we can say that the addition of two matrices satisfying the (graded) QYBE does not need to satisfy the (graded) QYBE. Also the equation (16) can be written of the form

$$R_q^1 T'_1 T'_2 = -T'_2 T'_1 R_q^2. \quad (22)$$

Let g_i and $\hat{T}_i$ ($i = 1, 2$) be graded in the same manner as $\hat{T}'_i$. It is clear that

$$\hat{T}'_1 \hat{T}'_2 = G_h \hat{T}_1 \hat{T}_2 G_h \quad \text{and} \quad \hat{T}'_2 \hat{T}'_1 = G_{-h} \hat{T}_2 \hat{T}_1 G_{-h}, \quad (23)$$

where

$$G_h = g_1 g_2^{-1}, \quad G_{-h} = G_h^{-1}. \quad (24)$$

Substituting (23) into (16) we arrive at the equation

$$G_h R_q G_h \hat{T}_1 \hat{T}_2 = -\hat{T}_2 \hat{T}_1 G_{-h} R_q G_{-h}. \quad (25)$$

If we define the R-matrix R_h as

$$R_h = \lim_{q \rightarrow 1} (G_h R_q G_h), \quad (26)$$

[after dividing by 2] we get the following R-matrix R_h

$$R_h = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -h & -1 & 0 & 0 \\ -h & 0 & -1 & 0 \\ 0 & h & -h & 1 \end{pmatrix} \quad (27)$$

which gives the relations (10) with the equation

$$R_h \hat{T}_1 \hat{T}_2 = -\hat{T}_2 \hat{T}_1 R_{-h}. \quad (28)$$

Note that although R_h does not satisfy the usual graded and ungraded YBE (19) it satisfies the equation

$$(R_{12} R_{13} R_{23})(h) = (R_{23} R_{13} R_{12})(-h), \quad (29)$$

or, equivalently,

$$R_{12} R_{13} R_{23} + R_{23} R_{13} R_{12} = 2I_8, \quad (30)$$

where I_8 is the 8x8 unit matrix and

$$R_{12} = R \otimes I_2, \quad \text{etc.}$$

Since

$$R_h^2 = I,$$

the R-matrix R_h has two eigenvalues ± 1 .

Acknowledgement

This work was supported in part by **T. B. T. A. K.** the Turkish Scientific and Technical Research Council.

References

- [1] V. G. Drinfeld, *Quantum groups*, in *Proc. IMS*, Berkeley, 1986;
N. Y. Reshetikhin, L. A. Takhtajan and L. D. Faddeev, *Leningrad Math. J.* **1**, 193 (1990).
- [2] E. E. Demidov, Yu. I. Manin, E. E. Mukhin and D. V. Zhdanovich, *Prog. Theor. Phys. Suppl.* **102**, 203 (1990);
H. Ewen, O. Ogievetsky and J. Wess, *Lett. Math. Phys.* **22**, 297 (1991);
S. Zakrzewski, *Lett. Math. Phys.* **22**, 287 (1991);
S. L. Woronowicz, *Rep. Math. Phys.* **30**, 259 (1991);
C. H. Ohn, *Lett. Math. Phys.* **25**, 85 (1992).
- [3] B. A. Kupershmidt, *J. Phys. A: Math. Gen.* **25**, L1239 (1992);
V. Karimipour, *Lett. Math. Phys.* **30**, 87 (1994);
A. Aghamohammadi, *Mod. Phys. Lett. A* **8**, 2607 (1993).
S. Celik and E. Hızal, *h-deformation of $Gr(2)$* , to appear in *J. Phys. A: Math. Gen.* **30** (1997), 4677-4680.
- [4] A. Aghamohammadi, M. Khorrami and A. Shariati, *J. Phys. A: Math. Gen.* **28**, L225 (1995).
- [5] S. Celik and S. A. Celik, *Balkan Phys. Lett.* **3**, 188 (1995).
- [6] Yu. I. Manin, *Commun. Math. Phys.* **123**, 163 (1989).
- [7] L. Dabrowski and P. Parashar, *h-deformation of $GL(1|1)$* to appear in *Lett. Math. Phys.*
- [8] S. Celik, *J. Math. Phys.* **37**, 3568 (1996).